

Tropical Jacobians in $\mathbb{R}^2$

Shuhei Yoshitomi

Abstract

In this paper, we will show that the topological type of the Jacobian variety of a reduced tropical plane curve depends only on the genus of the curve.

1 Introduction

This paper is a continuation of [7].

Definition. The *genus* of a tropical curve C in $\mathbb{R}^2$ is the number of lattice points in the Newton complex $\text{Newt}(C)$ which lies in the interior of the Newton polygon $\Delta(C)$. (Hence if C is reduced, the genus of C equals the rank of $H_1(C, \mathbb{Z})$.)

Definition. The *Jacobian variety* $\text{Jac}(C)$ is the residue group of $\text{Div}^0(C)$ by linear equivalence. Divisors D, D' are *linearly equivalent*, $D \sim D'$, if there are tropical curves L, L' such that $\Delta(L) = \Delta(L')$ and $D - D' = C \cdot L - C \cdot L'$.

Theorem 1.1. *Let C be a reduced tropical curve of genus g in $\mathbb{R}^2$. Then $\text{Jac}(C)$ is isomorphic to the g -dimensional real torus T^g .*

See [7] section 2 for preliminaries.

2 Proof of the theorem

Lemma 2.1 ([7], Lemma 3.2). *Let E be an edge of C , and let P, P', Q, Q' be points of E such that $\overrightarrow{PP'} = \overrightarrow{QQ'}$. Then $P' - P \sim Q' - Q$.*

Lemma 2.2 ([7], Lemma 3.4). *Let E be a ray of C . Then all points of E are linearly equivalent.*

Let $\mathcal{P}(\mathbb{R}^2)$ be the space of all tropical curves (See [7], section 4).

Lemma 2.3 ([7], Corollary 4.2). *Tropical curves $L, L' \in \mathcal{P}(\mathbb{R}^2)$ lie in the same connected component if and only if $\Delta(L) = \Delta(L')$.*

We consider the kernel of the homomorphism $\text{Div}^0(C) \rightarrow \text{Jac}(C)$. Lemma 2.1 says some subgroup $H \subset \text{Div}^0(C)$ is contained in this kernel. H is the subgroup generated by all $(P' - P) - (Q' - Q)$ which satisfies the assumption of Lemma 2.1.

Let $E_1, \dots, E_N$ be all finite edges of C . Let V_i be one of vertices of E_i . For each i , we have a homomorphism

$$\begin{aligned} \pi_i: \mathbb{R} &\longrightarrow \text{Div}^0(C)/H \\ \delta &\longmapsto P - V_i \quad (\delta > 0 \text{ is small enough}), \end{aligned}$$

where $P \in E_i$ is the point such that the lattice length of $[V_i, P]$ equals δ . Hence we have a homomorphism

$$\pi: \mathbb{R}^N \longrightarrow \text{Div}^0(C)/H.$$

Let $F = \text{Ker } \pi$. The composite map

$$\mathbb{R}^N/F \longrightarrow \text{Div}^0(C)/H \longrightarrow \text{Jac}(C)$$

is surjective because of Lemma 2.2.

We fix a vertex $\mathcal{O} \in C$. Let $\lambda: C \rightarrow \mathbb{R}^N/F$ be the map such that

i) $\pi(\lambda(P)) = P - \mathcal{O}$ for $P \in E_i$,

ii) every ray of C is contracted by λ .

Let $\tilde{\lambda}_i: E_i \rightarrow \mathbb{R}^N$ be a lift of $\lambda|_{E_i}: E_i \rightarrow \mathbb{R}^N/F$. Namely,

$$\tilde{\lambda}_i(P) = \tilde{\lambda}_i(V_i) + \delta e_i,$$

where δ is the lattice length of $[V_i, P]$.

Lemma 2.4. *F is a free abelian group of rank g .*

Proof. Let $\Lambda_1, \dots, \Lambda_g$ be cycles generating $H_1(C, \mathbb{Z})$. Let $V_{j1}, \dots, V_{js_j}$ be all vertices of Λ_j , and take $V_{j0} = V_{js_j}$. Let $a_{jk} \in \mathbb{R}^N$ be the vector such that $\pi(a_{jk}) = V_{jk} - V_{j,k-1}$. Let F' be the abelian group generated by $a_1, \dots, a_g$, where $a_j = \sum_k a_{jk}$. The map $\lambda: C \rightarrow \mathbb{R}^N/F$ is factored into a map

$$\lambda': C \longrightarrow \mathbb{R}^N/F'$$

and the homomorphism $\mathbb{R}^N/F' \rightarrow \mathbb{R}^N/F$. λ' induces a homomorphism $\lambda'': \text{Div}^0(C)/H \rightarrow \mathbb{R}^N/F'$. The homomorphism $\mathbb{R}^N \rightarrow \mathbb{R}^N/F'$ is factored into π and λ'' , so we have $F = F'$. $\square$

For a tropical curve $L \in \mathcal{P}(\mathbb{R}^2)$, we define $\sigma(L) \in \mathbb{R}^N/F$ as follows.

$$C \cdot L = P_1 + \cdots + P_r,$$

$$\sigma(L) = \lambda(P_1) + \cdots + \lambda(P_r).$$

The map

$$\sigma: \mathcal{P}(\mathbb{R}^2) \longrightarrow \mathbb{R}^N/F$$

is continuous by definition of the stable intersection.

Given a continuous path $\gamma: [0, 1] \rightarrow \mathcal{P}(\mathbb{R}^2)$, let $\tilde{\sigma}_\gamma: [0, 1] \rightarrow \mathbb{R}^N$ be a lift of $\sigma|_\gamma: [0, 1] \rightarrow \mathbb{R}^N/F$, and let $v(\gamma) = \tilde{\sigma}_\gamma(1) - \tilde{\sigma}_\gamma(0)$. Let $W \subset \mathbb{R}^N$ be the set of all $v(\gamma)$.

Lemma 2.5. *W is a linear subspace in $\mathbb{R}^N$.*

Proof. Recall that each component of $\mathcal{P}(\mathbb{R}^2)$ is a closed cone in an affine space. Given a Newton complex $\mathcal{N}$, let

$$\mathcal{P}(\mathcal{N}, C) = \{L \in \mathcal{P}(\mathbb{R}^2) \mid \text{Newt}(C \cup L) = \mathcal{N}\}.$$

Every element of W is written as $v(\gamma_1) + \cdots + v(\gamma_s)$ for linear paths

$$\gamma_j: [0, 1] \rightarrow \overline{\mathcal{P}(\mathcal{N}_j, C)}$$

and Newton complexes $\mathcal{N}_1, \dots, \mathcal{N}_s$. For $0 \leq t \leq 1$, we have

$$t(v(\gamma_1) + \cdots + v(\gamma_s)) = v(\gamma_{1,t}) + \cdots + v(\gamma_{s,t}),$$

where $\gamma_{j,t}: [0, t] \rightarrow \overline{\mathcal{P}(\mathcal{N}_j, C)}$ is a restriction of γ_j . Finally, $v(\gamma) + v(\gamma') \in W$, because $\sigma(L \cup L') = \sigma(L) + \sigma(L')$. $\square$

We have an isomorphism

$$\mathbb{R}^N/(F + W) \cong \text{Jac}(C)$$

from Lemma 2.3.

Note that $\mathbb{R}^N/W$ is a vector space generated by $e_1, \dots, e_N$. Theorem 1.1 follows from the following lemma.

Lemma 2.6. Suppose that $\mathcal{X} := \overline{C \setminus (E_1 \cup \dots \cup E_g)}$ is a maximal tree in C . Then $\langle e_1, \dots, e_g \rangle$ is a basis of $\mathbb{R}^N/W$ over $\mathbb{R}$.

Proof of that $\langle e_1, \dots, e_g \rangle$ is a generator. For $g+1 \leq j \leq N$, there is a piecewise linear curve $M \subset \mathbb{R}^2$ containing two rays such that $\mathcal{X} \cap M$ is a single point of E_j . The following lemma says that $e_j \in \mathbb{Q}\langle e_1, \dots, e_g \rangle$ in $\mathbb{R}^N/W$. $\square$

Lemma 2.7. Let S, T be rays in $\mathbb{R}^2$ such that

$$S \cap C = \emptyset, \quad T \cap C = \emptyset.$$

Let $P_0, \dots, P_r, P'_0, \dots, P'_r \in \mathbb{R}^2$ be rational points such that

$$P_0, P'_0 \in S, \quad P_r, P'_r \in T,$$

$$[P_i, P'_i] \cap C = \emptyset,$$

$$\overrightarrow{P_{i-1}P_i} \text{ and } \overrightarrow{P'_{i-1}P'_i} \text{ have the same direction.}$$

Then there are integers $m_1, \dots, m_r > 0$ such that

$$\sum_{i=1}^r m_i (C \cdot [P_{i-1}, P_i]) \sim \sum_{i=1}^r m_i (C \cdot [P'_{i-1}, P'_i]).$$

Proof. Let L be the tropical curve with vertex P_i , consisting of three rays L_0, L_1, L_2 such that

$$P_{i-1} \in L_1, \quad P_{i+1} \in L_2,$$

$$[P_i, P'_i] \text{ is parallel to } L_0.$$

Let $w_i = \text{wt}(L_1)$, $v_i = \text{wt}(L_2)$. Move L by $\overrightarrow{P_iP'_i}$, and denote it by L' . Comparing $C \cdot L$ and $C \cdot L'$, we have

$$C \cdot (w_i[P_i, P_{i-1}, \infty) \cup v_i[P_i, P_{i+1}, \infty)) \sim C \cdot (w_i[P'_i, P'_{i-1}, \infty) \cup v_i[P'_i, P'_{i+1}, \infty))$$

for $1 \leq i \leq r-1$. Similarly,

$$C \cdot (w_0[P_0, S] \cup v_0[P_0, P_1, \infty)) \sim C \cdot (w_0[P'_0, S] \cup v_0[P'_0, P'_1, \infty)),$$

$$C \cdot (w_r[P_r, P_{r-1}, \infty) \cup v_r[P_r, T]) \sim C \cdot (w_r[P'_r, P'_{r-1}, \infty) \cup v_r[P'_r, T]).$$

Taking a suitable linear combination and canceling lines, we have the statement. $\square$

Proof of that $\langle e_1, \dots, e_g \rangle$ is linearly independent. Suppose

$$a_1 e_1 + \dots + a_g e_g = v(\gamma),$$

$$\gamma: [0, 1] \longrightarrow \mathcal{P}(\mathbb{R}^2).$$

For $1 \leq j \leq g$, there is a piecewise linear closed curve Λ in $\mathcal{X} \cup E_j$. Let $E_{i(0)}, \dots, E_{i(s)}$ be all edges of Λ , and take $E_{i(0)} = E_j$. Let $\varepsilon_k \in \{1, -1\}$ be the sign of $E_{i(k)}$ with respect to the positive direction of S^1 : The sign is positive if the coordinate of $\mathbb{R}^N$ is increasing. Let $h: \mathbb{R}^N \rightarrow \mathbb{R}$ be the linear map defined by

$$h = \varepsilon_0 \check{e}_{i(0)} + \dots + \varepsilon_s \check{e}_{i(s)},$$

where $\check{e}_i$ denotes the dual basis. Then we have

$$h(\tilde{\sigma}_\gamma(1)) - h(\tilde{\sigma}_\gamma(0)) = h(v(\gamma)) = \varepsilon_0 a_j.$$

To complete the proof, we show that the map $h \circ \tilde{\sigma}_\gamma: [0, 1] \rightarrow \mathbb{R}$ is constant for any continuous path γ .

We may assume that C intersects $L_t := \gamma(t)$ transversely for any t . There are points $P_{kmt} \in E_{i(k)}$, edges $L_{kmt} \subset L_t$, and vectors $u_{km} \in \mathbb{R}^2$ such that

- i) $E_{i(k)} \cdot L_t = \sum_m P_{kmt}$,
- ii) $E_{i(k)} \cap L_{kmt} = P_{kmt}$,
- iii) u_{km} is the weighted primitive vector of L_{kmt} starting at the vertex inside Λ , divided by $\mu_{P_{kmt}}$.

For L_0 and L_1 , we have the moment condition inside Λ :

$$\sum_{k,m} \text{moment}(u_{km}, P_{km0}) = 0,$$

$$\sum_{k,m} \text{moment}(u_{km}, P_{km1}) = 0.$$

From these,

$$\sum_{k,m} \left(\overrightarrow{P_{km0} P_{km1}} \times u_{km} \right) = 0.$$

Let u_k be the primitive vector of $E_{i(k)}$ starting at $V_{i(k)}$. Since $u_k \times u_{km} = -\varepsilon_k$, we have

$$\sum_{k,m} h \left(\tilde{\lambda}_{i(k)}(P_{km0}) - \tilde{\lambda}_{i(k)}(P_{km1}) \right) = 0.$$

Thus $h(\tilde{\sigma}_\gamma(0)) = h(\tilde{\sigma}_\gamma(1))$. □

References

- [1] T. Bogart, A. Jensen, D. Speyer, B. Sturmfels, R. Thomas. Computing tropical varieties. Preprint, arXiv:math.AG/0507563.
- [2] A. Gathmann. Tropical algebraic geometry. Preprint, arXiv:math.AG/0601322.
- [3] J. Richter-Gebert, B. Sturmfels, and T. Theobald. First Steps in Tropical Geometry. Preprint, arXiv:math.AG/0306366.
- [4] Z. Izhakian. Tropical Varieties, Ideals and An Algebraic Nullstellensatz. Preprint, arXiv:math.AC/0511059.
- [5] G. Mikhalkin. Enumerative tropical algebraic geometry in $\mathbb{R}^2$. Preprint, arXiv:math.AG/0312530.
- [6] M. D. Vigeland. The group law on a tropical elliptic curve. Preprint, arXiv:math.AG/0411485.
- [7] S. Yoshitomi. Jacobian varieties of reduced tropical curves. Preprint, arXiv:math.AG/0612810.